



(19)

Europäisches Patentamt

European Patent Office

Office européen des brevets

(11)

EP 0 924 297 A1

EUROPEAN PATENT APPLICATION

(51) Int Cl.: C12N 15/12, C12N 15/85, C12N 5/10, C12Q 1/68, C07K 14/705, C07K 16/28, G01N 33/50, A61K 48/00

(43) Date of publication: 23.06.1999 Bulletin 1999/25

(21) Application number: 98302537.0

(22) Date of filing: 01.04.1998

(84) Designated Contracting States: AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU MC NL PT SE Designated Extension States: AL LT LV MK RO SI Priority: 02.04.1997 US 41886 P

(71) Applicant: SMITHKLINE BEECHAM CORPORATION

Philadelphia Pennsylvania 19103 (US)

(54) Sialoadhesin family member-2 (saf-2)

(57) The invention relates to a new member of the Sialoadhesin gene family, SAF-2 polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing SAF-2 polypeptides and polynucleotides in therapy, and diagnostic assays for such.

Description

[0001] This application claims the benefit of U.S. Provisional Application No. 60/041,886, filed April 2, 1997, whose contents are incorporated herein by reference.

Field of the invention

[0002] This invention relates to newly identified polypeptides and polynucleotides encoding such polypeptides, to their use in therapy and in identifying compounds which may be agonists, antagonists and/or inhibitors which are potentially useful in therapy, and to production of such polypeptides and polynucleotides.

Background of the invention

[0003] The drug discovery process is currently undergoing a fundamental revolution as it embraces functional genomics, that is, high throughput genome- or gene-based biology. This approach is rapidly superseding earlier approaches based on positional cloning. A phenotype, that is a biological function or genetic disease, would be identified and this would then be tracked back to the responsible gene, based on its genetic map position.

[0004] Functional genomics relies heavily on the various tools of bioinformatics to identify gene sequences of potential interest from the many molecular biology databases now available. There is a continuing need to identify and characterise further genes and their related polypeptides/proteins, as targets for drug discovery.

Summary of the invention

[0005] The present invention relates to SAF-2, in particular SAF-2 polypeptides and SAF-2 polynucleotides, recombinant materials and methods for their production. In another aspect, the invention relates to methods for using such polypeptides and polynucleotides, including the treatment of cancer, inflammation, autoimmune, allergy, asthma, rheumatoid arthritis, CNS inflammation, cerebellar degeneration, Alzheimer's disease, Parkinson's disease, multiple sclerosis, amyotrophic lateral sclerosis, head injury damage, and other neurological abnormalities, septic shock, sepsis, stroke, osteoporosis, osteoarthritis, ischemia reperfusion injury, cardiovascular disease, kidney disease, liver disease, ischemic injury, myocardial infarction, hypotension, hypertension, AIDS, myelodysplastic syndromes and other hematologic abnormalities, aplastic anemia, male pattern baldness, and bacterial, fungal, protozoan and viral infections, hereinafter referred to as "the Diseases", amongst others. In a further aspect, the invention relates to methods for identifying agonists and antagonists/inhibitors using the materials provided by the invention, and treating conditions associated with SAF-2 imbalance with the identified compounds. In a still further aspect, the invention relates to diagnostic assays for detecting diseases associated with inappropriate SAF-2 activity or levels.

Description of the invention

[0006] In a first aspect, the present invention relates to SAF-2 polypeptides. Such peptides include isolated polypeptides comprising an amino acid sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, that of SEQ ID NO:2 over the entire length of SEQ ID NO:2. Such polypeptides include those comprising the amino acid of SEQ ID NO:2.

[0007] Further peptides of the present invention include isolated polypeptides in which the amino acid sequence has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2. Such polypeptides include the polypeptide of SEQ ID NO:2.

[0008] Further peptides of the present invention include isolated polypeptides encoded by a polynucleotide comprising the sequence contained in SEQ ID NO:1.

[0009] Polypeptides of the present invention are believed to be members of the sialoadhesin family of polypeptides. They are therefore of interest because the sialoadhesin family of proteins, including sialoadhesin, CD33, CD22 and myelin-associated glycoprotein (MAG), are utilized as cellular interaction molecules. These proteins bind specific carbohydrates in a static acid dependent manner on target cells. The extracellular domain is made up of various numbers of immunoglobulin-like domains of the V-like and C2-like subtypes, and the intracellular portion has no homology to any known signalling motifs. Sialoadhesin expression is restricted to macrophages; it has 17 Ig-like domains and the specific recognition sequence on target cells is Neu5Ac α 2,3Gal β 13GalNAc. Known target cells include developing myeloid cells in the bone marrow and lymphocytes in the spleen and lymph node (Crocker, P. R., et al. EMBO J, 1994, 13:4490-4503). CD22 is expressed only on B cells and has α and β isoforms with 5 and 7 Ig-like domains, respectively.

CD22 is known to bind T cells, B cells, monocytes, granulocytes and erythrocytes by recognizing Neu5Ac2,6Gal1,4Glc(Nac) in N-linked glycans (Crocke, R. J., et al. EMBO J. 1994, 13:4490-4503; Stamenkovic, I. and Seed, B. Nature, 1990, 345:74-77; Wilson, G. L., et al. J Exp Med. 1991, 173:137-146). Myelin-associated glycoprotein (MAG) is expressed by Schwann cells in the peripheral nervous system and oligodendrocytes in the central nervous system and is thought to participate in the cell adhesion to axons. MAG has two alternatively spliced variants, large MAG (L-MAG) and small MAG (S-MAG), which are expressed either during embryonic development or in adult, respectively. The alternative splicing results in the expression of the same extracellular domains but distinct intracellular domains (Pedraza, L. et al. JCB, 1990, 111:2651-2661).

[0010] CD33 is most relevant to SAF-2 because they are the most closely related of all the family members. CD33 is normally expressed on the developing myelomonocytic lineage. It is absent on early stem cells but is present on colony-forming units for granulocytes, erythrocytes, and megakaryocytes (CFU-GEMM) and progenitors of granulocytes and mononuclear phagocytes (CFU-GM). It is down-regulated by mature granulocytes and retained by large MAG (L-MAG) and macrophages (Andrews, R. G., et al. Blood, 1983, 62:124; Griffin, J. D., et al. Leuk Res 1984, 8:521). CD33 has two Ig-like domains and has an affinity for target cells expressing NeuAc2,3Gal in N- and O-linked glycans. It maps to chromosome 19q13.1-13.3, and is, therefore, closely linked in the genome with MAG and CD22 (Freeman, S. D., et al. Blood, 1995, 85:2005-2012).

[0011] CD33 has also been found to be expressed on about 85% of leukemic myeloblasts in patients with acute myelogenous leukemia (AML) and is frequently used to differentiate AML from acute lymphoblastic leukemia (ALL). Monoclonal antibodies to CD33 have been used therapeutically to purge residual myeloblasts from autologous bone marrow grafts *ex vivo* for the treatment of AML (Robertson, M. J., et al. Blood, 1992, 79:2229-2236). More recently, humanized monoclonal antibodies to CD33 have undergone evaluation *in vivo* for the treatment of AML (Caron, P. C., et al. Blood, 1994, 83:1760-1768). These properties are hereinafter referred to as "SAF-2 activity" or "SAF-2 polypeptide activity" or "biological activity of SAF-2". Also included amongst these activities are antigenic and immunogenic activities of said SAF-2 polypeptides, in particular the antigenic and immunogenic activities of the polypeptide of SEQ ID NO:2. Preferably a polypeptide of the present invention exhibits at least one biological activity of SAF-2.

[0012] The polypeptides of the present invention may be in the form of the "mature" protein or may be a part of a larger protein such as a fusion protein. It is often advantageous to include an additional amino acid sequence which contains secretory or leader sequences, pro-sequences, sequences which aid in purification such as multiple histidine residues, or an additional sequence for stability during recombinant production.

[0013] The present invention also includes variants of the aforementioned polypeptides, that is polypeptides that vary from the references by conservative amino acid substitutions, whereby a residue is substituted by another with like characteristics. Typical such substitutions are among Ala, Val, Leu and Ile; among Ser and Thr; among the acidic residues Asp and Glu; among Asn and Gln; and among the basic residues Lys and Arg; or aromatic residues Phe and Tyr. Particularly preferred are variants in which several, 5-10, 1-5, 1-3, 1-2 or 1 amino acids are substituted, deleted, or added in any combination.

[0014] Polypeptides of the present invention can be prepared in any suitable manner. Such polypeptides include isolated naturally occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art.

[0015] In a further aspect, the present invention relates to SAF-2 polynucleotides. Such polynucleotides include isolated polynucleotides comprising a nucleotide sequence encoding a polypeptide which has at least 70% identity, yet more preferably at least 80% identity, to the amino acid sequence of SEQ ID NO:2, over the entire length of SEQ ID NO:2. In this regard, polypeptides which have at least 97% identity are highly preferred, whilst those with at least 98-99% identity are most highly preferred, and those with at least 99% identity are most highly preferred. Such polynucleotides include a polynucleotide comprising the nucleotide sequence contained in SEQ ID NO:1 encoding the polypeptide of SEQ ID NO:2.

[0016] Further polynucleotides of the present invention include isolated polynucleotides comprising a nucleotide sequence that has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, to a nucleotide sequence encoding a polypeptide of SEQ ID NO:2, over the entire coding region. In this regard, polynucleotides which have at least 97% identity are highly preferred, whilst those with at least 98-99% identity are more highly preferred, and those with at least 99% identity are most highly preferred.

[0017] Further polynucleotides of the present invention include isolated polynucleotides comprising a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, to SEQ ID NO:1 over the entire length of SEQ ID NO:1. In this regard, polynucleotides which have at least 97% identity are highly preferred, whilst those with at least 98-99% identity are more highly preferred, and those with at least 99% identity are most highly preferred. Such polynucleotides include a polynucleotide comprising the polynucleotide of SEQ ID NO:1 as well as the polynucleotide of SEQ ID NO:1.

[0018] The invention also provides polynucleotides which are complementary to all the above described polynucle-

olides.

[0019] The nucleotide sequence of SEQ ID NO:1 shows homology with CD33 (Simmons, D., and Seed, B., J1 141: 2797-2800, 1988). The nucleotide sequence encoding the polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence contained in SEQ ID NO:1 or it may be a sequence other than the one contained in SEQ ID NO:1, which, as a result of the redundancy (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2. The polypeptide of the SEQ ID NO:2 is structurally related to other proteins of the staphylococcal family, having homology and/or structural similarity with CD33 (Simmons, D., and Seed, B., J1 141: 2797-2800, 1988).

[0020] Preferred polypeptides and polynucleotides of the present invention are expected to have, *inter alia*, similar biological functions/properties to their homologous polypeptides and polynucleotides. Furthermore, preferred polypeptides and polynucleotides of the present invention have at least one SAF-2 activity.

[0021] The present invention also relates to partial or other polynucleotide and polypeptide sequences which were first identified prior to the determination of the corresponding full length sequences of SEQ ID NO:1 and SEQ ID NO:2. Accordingly, in a further aspect, the present invention provides for an isolated polynucleotide comprising:

(a) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity to SEQ ID NO:3 over the entire length of SEQ ID NO:3;

(b) a nucleotide sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity, to SEQ ID NO:3 over the entire length of SEQ ID NO:3;

(c) the polynucleotide of SEQ ID NO:3; or

(d) a nucleotide sequence encoding a polypeptide which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, even more preferably at least 97-99% identity, to the amino acid sequence of SEQ ID NO:4 over the entire length of SEQ ID NO:4;

(a) comprises an amino acid sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to that of SEQ ID NO:4 over the entire length of SEQ ID NO:4;

(b) has an amino acid sequence which is at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to the amino acid sequence of SEQ ID NO:4 over the entire length of SEQ ID NO:4;

(c) comprises the amino acid of SEQ ID NO:4; and

(d) is the polypeptide of SEQ ID NO:4;

as well as the polynucleotide of SEQ ID NO:3.

[0023] The present invention further provides for a polypeptide which:

(a) comprises an amino acid sequence which has at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to that of SEQ ID NO:4 over the entire length of SEQ ID NO:4;

(b) has an amino acid sequence which is at least 70% identity, preferably at least 80% identity, more preferably at least 90% identity, yet more preferably at least 95% identity, most preferably at least 97-99% identity, to the amino acid sequence of SEQ ID NO:4 over the entire length of SEQ ID NO:4;

(c) comprises the amino acid of SEQ ID NO:4; and

(d) is the polypeptide of SEQ ID NO:4;

as well as polypeptides encoded by a polynucleotide comprising the sequence contained in SEQ ID NO:3.

[0024] The nucleotide sequence of SEQ ID NO:3 and the peptide sequence encoded thereby are derived from EST (Expressed Sequence Tag) sequences. It is recognised by those skilled in the art that there will inevitably be some nucleotide sequence reading errors in EST sequences (see Adams, M.D. *et al.* Nature 377 (supp) 3, 1995). Accordingly, the nucleotide sequence of SEQ ID NO:3 and the peptide sequence encoded therefrom are therefore subject to the same inherent limitations in sequence accuracy. Furthermore, the peptide sequence encoded by SEQ ID NO:3 comprises a region of identity or close homology and/or close structural similarity (for example a conservative amino acid difference) with the closest homologous or structurally similar protein.

[0025] Polynucleotides of the present invention may be obtained, using standard cloning and screening techniques, from a cDNA library derived from mRNA in cells of human eosinophils, using the expressed sequence tag (EST) analysis (Adams, M.D. *et al.* Science (1991) 252:1651-1656; Adams, M.D. *et al.* Nature, (1992) 355:632-634; Adams, M.D. *et al.* Nature (1995) 377 Supp:3-174). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

[0026] When polynucleotides of the present invention are used for the recombinant production of polypeptides of the present invention, the polynucleotide may include the coding sequence for the mature polypeptide, by itself, or the coding sequence for the mature polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro-protein sequence, or other fusion peptide portions. For example, a marker sequence which facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pCE vector

(Clagen, Inc.) and described in Gentz *et al.*, *Proc Natl Acad Sci USA* (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

[0027] Further embodiments of the present invention include polynucleotides encoding polypeptide variants which comprise the amino acid sequence of SEQ ID NO:2 and in which several, for instance from 5 to 10, 1 to 5, 1 to 3, 1 to 2 or 1, amino acid residues are substituted, deleted or added, in any combination.

[0028] Polynucleotides which are identical or sufficiently identical to a nucleotide sequence contained in SEQ ID NO:1, may be used as hybridization probes for cDNA and genomic DNA or as primers for a nucleic acid amplification (PCR) reaction, to isolate full-length cDNAs and genomic clones encoding polypeptides of the present invention and to isolate cDNA and genomic clones of other genes (including genes encoding homologs and orthologs from species other than human) that have a high sequence similarity to SEQ ID NO:1. Typically these nucleotide sequences are 70% identical, preferably 80% identical, more preferably 90% identical, most preferably 95% identical to that of the reference. The probes or primers will generally comprise at least 15 nucleotides, preferably, at least 30 nucleotides and may have at least 50 nucleotides. Particularly preferred probes will have between 30 and 50 nucleotides.

[0029] A polynucleotide encoding a polypeptide of the present invention, including homologs and orthologs from species other than human, may be obtained by a process which comprises the steps of screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or a fragment thereof, and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to the skilled artisan. Preferred stringent hybridization conditions include overnight incubation at 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA; followed by washing the filters in 0.1x SSC at about 65°C. Thus the present invention also includes polynucleotides obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or a fragment thereof.

[0030] The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide is cut short at the 5' end of the cDNA. This is a consequence of reverse transcriptase, an enzyme with inherently low processivity (a measure of the ability of the enzyme to remain attached to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during 1st strand cDNA synthesis.

[0031] There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extended short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) (see, for example, Frohman *et al.*, PNAS USA 85, 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon™ technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon™ technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated onto each end. Nucleic acid amplification (PCR) is then carried out to amplify the 'missing' 5' end of the cDNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using 'nested' primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the known gene sequence). The products of this reaction can then be analysed by DNA sequencing and a full-length cDNA constructed either by joining the product directly to the existing cDNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

[0032] Recombinant polypeptides of the present invention may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems which comprise a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

[0033] For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Introduction of polynucleotides into host cells can be effected by methods described in many standard laboratory manuals, such as Davis *et al.*, *Basic Methods in Molecular Biology* (1986) and Sambrook *et al.*, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989). Preferred such methods include, for instance, calcium phosphate transfection, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

[0034] Representative examples of appropriate hosts include bacterial cells, such as *Streptococcus*, *Staphylococcus*, *coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

[0035] A great variety of expression systems can be used, for instance, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector which is able to maintain, propagate or express a polynucleotide to produce a polypeptide in a host may be used. The appropriate nucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, *MOLECULAR CLONING, A LABORATORY MANUAL* (*supra*). Appropriate secretion signals may be incorporated into the desired polypeptide to allow secretion of the translated protein into the lumen of the endoplasmic reticulum, the periplasmic space or the extracellular environment. These signals may be endogenous to the polypeptide or they may be heterologous signals.

[0036] If a polypeptide of the present invention is to be expressed for use in screening assays, it is generally preferred that the polypeptide be produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide. If produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

[0037] Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxyapatite chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during isolation and/or purification.

[0038] This invention also relates to the use of polynucleotides of the present invention as diagnostic reagents. Detection of a mutated form of the gene characterised by the polynucleotide of SEQ ID NO:1 which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from under-expression, over-expression or altered expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques.

[0039] Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or may be amplified enzymatically by using PCR or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by hybridizing amplified DNA to labeled SAF-2 nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence differences may also be detected by alterations in electrophoretic mobility of DNA fragments in gels, with or without denaturing agents, or by direct DNA sequencing (see, e.g., Myers *et al.*, *Science* (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton *et al.*, *Proc Natl Acad Sci USA* (1985) 85:4397-4401). In another embodiment, an array of oligonucleotides probes comprising SAF-2 nucleotide sequence or fragments thereof can be constructed to conduct efficient screening of e.g., genetic mutations. Array technology methods are well known and have general applicability and can be used to address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability (see for example: M.Chee *et al.*, *Science*, Vol 274, pp 610-613 (1996)).

[0040] The diagnostic assays offer a process for diagnosing or determining a susceptibility to the Diseases through detection of mutation in the SAF-2 gene by the methods described. In addition, such diseases may be diagnosed by methods comprising determining from a sample derived from a subject an abnormally decreased or increased level of polypeptide or mRNA. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays and flow cytometric analysis.

[0041] Thus in another aspect, the present invention relates to a diagnostic kit which comprises:

(a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 1, or a fragment thereof;

(b) a nucleotide sequence complementary to that of (a);

(c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or a fragment thereof; or

(d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO.2.

[0042] It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, particularly cancer, inflammation, autoimmunity, allergy, asthma, rheumatoid arthritis, CNS inflammation, cerebellar degeneration, Alzheimer's disease, Parkinson's disease, multiple sclerosis, amyotrophic lateral sclerosis, head injury damage, and other neurological abnormalities; septic shock, sepsis, stroke, osteoporosis, osteoarthritis, ischemia reperfusion injury, cardiovascular disease, kidney disease, liver disease, ischemic injury, myocardial infarction, hypotension, hypertension, AIDS, myelodysplastic syndromes and other hematologic abnormalities, aplastic anemia, male pattern baldness, and bacterial, fungal, protozoan and viral infections, amongst others.

[0043] The nucleotide sequences of the present invention are also valuable for chromosome identification. The sequence is specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian inheritance in Man (available on-line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (coinheritance of physically adjacent genes). The differences in the cDNA or genomic sequence between affected and unaffected individuals can also be determined. If a mutation is observed in some or all of the affected individuals but not in any normal individuals, then the mutation is likely to be the causative agent of the disease. The gene of the present invention maps to human chromosome 19q13.1-13.3.

[0045] The polypeptides of the invention or their fragments or analogs thereof, or cells expressing them, can also be used as immunogens to produce antibodies immunospecific for polypeptides of the present invention. The term "immunospecific" means that the antibodies have substantially greater affinity for the polypeptides of the invention than their affinity for other related polypeptides in the prior art.

[0046] Antibodies generated against polypeptides of the present invention may be obtained by administering the polypeptides or epitope-bearing fragments, analogs or cells to an animal, preferably a non-human animal, using routine protocols. For preparation of monoclonal antibodies, any technique which provides antibodies produced by continuous cell line cultures can be used. Examples include the hybridoma technique (Kohler, G. and Milstein, C., *Nature* (1975) 256:495-497), the trioma technique, the human B-cell hybridoma technique (Kozbor *et al.*, *Immunology Today* (1983) 4:72) and the EBV-hybridoma technique (Cole *et al.*, *MONOCLONAL ANTIBODIES AND CANCER THERAPY*, pp. 77-96, Alan R. Liss, Inc., 1985).

[0047] Techniques for the production of single chain antibodies, such as those described in U.S. Patent No. 4,946,778, can also be adapted to produce single chain antibodies to polypeptides of this invention. Also, transgenic mice, or other organisms, including other mammals, may be used to express humanized antibodies.

[0048] The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography.

[0049] Antibodies against polypeptides of the present invention may also be employed to treat the diseases, amongst others. Antibodies against SAF-2 polypeptides may also be employed to subcharacterize cell populations during hematopoietic development, as a diagnostic marker to distinguish between different forms of cancer, to purge bone marrow *ex vivo* of cancer cells expressing SAF-2, as a tool to aid in the *ex vivo* expansion (proliferation and/or differentiation) of hematopoietic progenitor cells expressing SAF-2, as a stimulus *in vivo* for stem cell mobilization into the periphery, and as an *in vivo* chemoprotective agent.

[0050] In a further aspect, the present invention relates to genetically engineered soluble fusion proteins comprising a polypeptide of the present invention, or a fragment thereof, and various portions of the constant regions of heavy or light chains of immunoglobulins or various subclasses (IgG, IgM, IgA, IgE). Preferred as an immunoglobulin is the constant part of the heavy chain of human IgG, particularly IgG1, where fusion takes place at the hinge region. In a particular embodiment, the Fc part can be removed simply by incorporation of a cleavage sequence which can be cleaved with blood clotting factor Xa. Furthermore, this invention relates to processes for the preparation of these fusion proteins by genetic engineering, and to the use thereof for drug screening, diagnosis and therapy. In another approach, soluble forms of SAF-2 polypeptides still capable of binding the ligand in competition with endogenous SAF-2 may be administered. Typical embodiments of such competitors comprise fragments of the SAF-2 polypeptide. One example is using the extracellular domain of SAF-2 fused to a human immunoglobulin Fc region which could then be employed to treat cancer, inflammation, autoimmunity and allergy, among others. SAF-2/Fc polypeptides may also be employed to purge bone marrow *ex vivo* of cancer cells expressing SAF-2 ligands, as a tool to aid in the *ex vivo* expansion (proliferation and/or differentiation) of hematopoietic progenitor cells expressing SAF-2 ligands, as a stimulus *in vivo* for stem cell mobilization into the periphery, and as an *in vivo* chemoprotective agent.

the invention also relates to polynucleotides encoding such fusion proteins. Examples of fusion protein technology can be found in International Patent Application Nos. WO94/29458 and WO94/22914.

[0051] Another aspect of the invention relates to a method for inducing an immunological response in a mammal which comprises inoculating the mammal with a polypeptide of the present invention, adequate to produce an antibody and/or T cell immune response to protect said animal from the Diseases hereinbefore mentioned, amongst others. Yet another aspect of the invention relates to a method of inducing immunological response in a mammal which comprises delivering a polypeptide of the present invention *via* a vector directing expression of the polynucleotide and coding for the polypeptide *in vivo* in order to induce such an immunological response to produce antibody to protect said animal from diseases.

[0052] A further aspect of the invention relates to an immunological/vaccine formulation (composition) which, when introduced into a mammalian host, induces an immunological response in that mammal to a polypeptide of the present invention wherein the composition comprises a polypeptide or polynucleotide of the present invention. The vaccine formulation may further comprise a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered parenterally (for instance, subcutaneous, intramuscular, intravenous, or intradermal injection). Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in-water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

[0053] Polypeptides of the present invention are responsible for many biological functions, including many disease states, in particular the Diseases hereinbefore mentioned. It is therefore desirable to devise screening methods to identify compounds which stimulate or which inhibit the function of the polypeptide. Accordingly, in a further aspect, the present invention provides for a method of screening compounds to identify those which stimulate or which inhibit the function of the polypeptide. In general, agonists or antagonists may be employed for therapeutic and prophylactic purposes for such Diseases as hereinbefore mentioned. Compounds may be identified from a variety of sources, for example, cells, cell-free preparations, chemical libraries, and natural product mixtures. Such agonists, antagonists or inhibitors so-identified may be natural or modified substrates, ligands, receptors, enzymes, etc., as the case may be, of the polypeptide; or may be structural or functional mimetics thereof (see Colligan *et al.*, *Current Protocols in Immunology* 1(2):Chapter 5 (1991)).

[0054] The screening method may simply measure the binding of a candidate compound to the polypeptide, or to cells or membranes bearing the polypeptide, or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve competition with a labeled competitor. Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells bearing the polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Constitutively active polypeptides may be employed in screening methods for inverse agonists or inhibitors, in the absence of an agonist or inhibitor, by testing whether the candidate compound results in inhibition of activation of the polypeptide. Further, the screening method may simply comprise the steps of mixing a candidate compound with a solution containing a polypeptide of the present invention, to form a mixture, measuring SAF-2 activity in the mixture, and comparing the SAF-2 activity of the mixture to a standard. Fusion proteins, such as those made from Fc portion and SAF-2 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists for the polypeptide of the present invention (see D. Bennett *et al.*, *J Mol Recognition*, 8:52-58 (1995); and K. Johanson *et al.*, *J Biol Chem*, 270(16):9459-9471 (1995)).

[0055] The polynucleotides, polypeptides and antibodies to the polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and of polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell-associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents which may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

[0056] The polypeptide may be used to identify membrane bound or soluble receptors, if any, through standard receptor binding techniques known in the art. These include, but are not limited to, ligand binding and crosslinking assays in which the polypeptide is labeled with a radioactive isotope (for instance, ¹²⁵I), chemically modified (for instance, biotinylated), or fused to a peptide sequence suitable for detection or purification, and incubated with a source of the putative receptor (cells, cell membranes, cell supernatants, tissue extracts, bodily fluids). Other methods include biophysical techniques such as surface plasmon resonance and spectroscopy. These screening methods may also be

used to identify agonists and antagonists of the polypeptide which compete with the binding of the polypeptide to its receptors, if any. Standard methods for conducting such assays are well understood in the art.

[0057] Examples of potential polypeptide antagonists include antibodies or, in some cases, oligonucleotides or proteins which are closely related to the ligands, substrates, receptors, enzymes, etc., as the case may be, of the polypeptide, e.g., a fragment of the ligands, substrates, receptors, enzymes, etc., or small molecules which bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

[0058] Thus, in another aspect, the present invention relates to a screening kit for identifying agonists, antagonists, ligands, receptors, substrates, enzymes, etc., for polypeptides of the present invention, or compounds which decrease or enhance the production of such polypeptides, which comprises:

- (a) a polypeptide of the present invention;
- (b) a recombinant cell expressing a polypeptide of the present invention;
- (c) a cell membrane expressing a polypeptide of the present invention;
- (d) antibody to a polypeptide of the present invention;

which polypeptide is preferably that of SEQ ID NO:2.

[0059] It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

[0060] It will be readily appreciated by the skilled artisan that a polypeptide of the present invention may also be used in a method for the structure-based design of an agonist, antagonist or inhibitor of the polypeptide, by:

- (a) determining in the first instance the three-dimensional structure of the polypeptide;
- (b) deducing the three-dimensional structure for the likely reactive or binding site(s) of an agonist, antagonist or inhibitor;
- (c) synthesizing candidate compounds that are predicted to bind to or react with the deduced binding or reactive site; and
- (d) testing whether the candidate compounds are indeed agonists, antagonists or inhibitors. It will be further appreciated that this will normally be an iterative process.

[0061] In a further aspect, the present invention provides methods of treating abnormal conditions such as, for instance, cancer, inflammation, autoimmunity, allergy, asthma, rheumatoid arthritis, CNS inflammation, cerebellar degeneration, Alzheimer's disease, Parkinson's disease, multiple sclerosis, amyotrophic lateral sclerosis, head injury damage, and other neurological abnormalities, septic shock, sepsis, stroke, osteoporosis, osteoarthritis, ischemia reperfusion injury, cardiovascular disease, kidney disease, liver disease, ischemic infarction, myocardial infarction, hypotension, hypereffusion, AIDS, myelodysplastic syndromes and other hematologic abnormalities, aplastic anemia, male pattern baldness, and bacterial, fungal, protozoan and viral infections, related to either an excess of, or an under-expression of, SAF-2 polypeptide activity.

[0062] If the activity of the polypeptide is in excess, several approaches are available. One approach comprises administering to a subject in need thereof an inhibitor compound (antagonist) as hereinabove described, optionally in combination with a pharmaceutically acceptable carrier, in an amount effective to inhibit the function of the polypeptide, such as, for example, by blocking the binding of ligands, substrates, receptors, enzymes, etc., or by inhibiting a second signal, and thereby alleviating the abnormal condition. In another approach, soluble forms of the polypeptides still capable of binding the ligand, substrate, enzymes, receptors, etc., in competition with endogenous polypeptide may be administered. Typical examples of such competitors include fragments of the SAF-2 polypeptide.

[0063] In still another approach, expression of the gene encoding endogenous SAF-2 polypeptide can be inhibited using expression blocking techniques. Known such techniques involve the use of antisense sequences, either internally generated or separately administered (see, for example, O'Connor, *J Neurochem* (1991) 56:560, in Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression, CRC Press, Boca Raton, FL (1988)). Alternatively, oligonucleotides which form triple helices with the gene can be supplied (see, for example, Lee et al., *Nucleic Acids Res* (1979) 6:3073; Cooney et al., *Science* (1988) 241:456; Dervan et al., *Science* (1991) 251:1360). These oligomers can be administered *per se* or the relevant oligomers can be expressed *in vivo*.

[0064] For treating abnormal conditions related to an under-expression of SAF-2 and its activity, several approaches are also available. One approach comprises administering to a subject a therapeutically effective amount of a compound which activates a polypeptide of the present invention, i.e., an agonist as described above, in combination with a pharmaceutically acceptable carrier, to thereby alleviate the abnormal condition. Alternatively, gene therapy may be employed to effect the endogenous production of SAF-2 by the relevant cells in the subject. For example, a polynucleotide of the invention may be engineered for expression in a replication defective retroviral vector, as discussed above. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding a polypeptide of the present invention such that the packaging

cell now produces infectious viral particles containing the gene of interest. These producer cells may be administered to a subject for engineering cells *in vivo* and expression of the polypeptide *in vivo*. For an overview of gene therapy, see Chapter 20, *Gene Therapy and other Molecular Genetic-based Therapeutic Approaches*, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996). Another approach is to administer a therapeutic amount of a polypeptide of the present invention in combination with a suitable pharmaceutical carrier.

[0065] In a further aspect, the present invention provides for pharmaceutical compositions comprising a therapeutically effective amount of a polypeptide, such as the soluble form of a polypeptide of the present invention, agonist/antagonist peptide or small molecule compound, in combination with a pharmaceutically acceptable carrier or excipient. Such carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The invention further relates to pharmaceutical packs and kits comprising one or more containers filled with one or more of the ingredients of the aforementioned compositions of the invention. Polypeptides and other compounds of the present invention may be employed alone or in conjunction with other compounds, such as therapeutic compounds.

[0066] The composition will be adapted to the route of administration, for instance by a systemic or an oral route. Preferred forms of systemic administration include injection, typically by intravenous injection. Other injection routes, such as subcutaneous, intramuscular, or intraperitoneal, can be used. Alternative means for systemic administration include transcutaneous and transdermal administration using penetrants such as bile salts or fusidic acids or other detergents. In addition, if a polypeptide or other compounds of the present invention can be formulated in an enteric or encapsulated formulation, oral administration may also be possible. Administration of these compounds may also be topical and/or localized, in the form of salves, pastes, gels, and the like.

[0067] The dosage range required depends on the choice of peptide or other compounds of the present invention, the route of administration, the nature of the formulation, the nature of the subject's condition, and the judgment of the attending practitioner. Suitable dosages, however, are in the range of 0.1-100 µg/kg of subject. Wide variations in the needed dosage, however, are to be expected in view of the variety of compounds available and the differing efficiencies of various routes of administration. For example, oral administration would be expected to require higher dosages than administration by intravenous injection. Variations in these dosage levels can be adjusted using standard empirical routines for optimization, as is well understood in the art.

[0068] Polypeptides used in treatment can also be generated endogenously in the subject, in treatment modalities often referred to as "gene therapy" as described above. Thus, for example, cells from a subject may be engineered with a polynucleotide, such as a DNA or RNA, to encode a polypeptide *ex vivo*, and for example, by the use of a retroviral plasmid vector. The cells are then introduced into the subject.

[0069] Polynucleotide sequences form a valuable information resource with which to identify further sequences of similar homology. This is most easily facilitated by storing the sequence in a computer readable medium and then using the stored data to search a sequence database using well known searching tools, such as GCG, Accelrys, and the like. The present invention provides for a computer readable medium having stored thereon a polynucleotide comprising the sequence of SEQ ID NO:1 and/or a polypeptide sequence encoded thereby.

[0070] The following definitions are provided to facilitate understanding of certain terms used frequently hereinbefore. "Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an Fab or other immunoglobulin expression library.

[0072] "Isolated" means altered "by the hand of man" from the natural state. If an "isolated" composition or substance occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living animal is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated," as the term is employed herein.

[0073] "Polynucleotide" generally refers to any polynucleotide or polynucleotide, which may be unmodified RNA or DNA or modified RNA or DNA. "Polynucleotides" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is a mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term "polynucleotide" also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

[0074] "Polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by

peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications may occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present to the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from post-translational natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphotylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination (see, for instance, PROTEINS - STRUCTURE AND MOLECULAR PROPERTIES, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993; Wolf, F., Post-translational Protein Modifications: Perspectives and Prospects, pgs. 1-12 in POSTTRANSLATIONAL COVALENT MODIFICATION OF PROTEINS, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol* (1990) 182:626-646 and Rattan et al., "Protein Synthesis: Post-translational Modifications and Aging", *Ann NY Acad Sci* (1992) 663:48-62).

[0075] "Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains essential properties. A typical variant of a polynucleotide differs in nucleotide sequence from another, reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from another, reference polypeptide. Generally, differences are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ in amino acid sequence by one or more substitutions, additions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. A variant of a polynucleotide or polypeptide may be a naturally occurring variant such as an allelic variant, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis.

[0076] "Identity" is a measure of the identity of nucleotide sequences or amino acid sequences. In general, the sequences are aligned so that the highest order match is obtained. "Identity" *per se* has an art-recognized meaning and can be calculated using published techniques (see, e.g., COMPUTATIONAL MOLECULAR BIOLOGY, Lesk, A. M., ed., Oxford University Press, New York, 1988; BIOCOMPUTING: INFORMATICS AND GENOME PROJECTS, Smith, D. W., ed., Academic Press, New York, 1993; COMPUTER ANALYSIS OF SEQUENCE DATA, PART I, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; SEQUENCE ANALYSIS IN MOLECULAR BIOLOGY, von Heinig, G., Academic Press, 1987; and SEQUENCE ANALYSIS PRIMER, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York, 1991). While there exist a number of methods to measure identity between two polynucleotide or polypeptide sequences, the term "identity" is well known to skilled artisans (Carillo, H., and Lipson, D., *SIAM J Applied Math* (1988) 48:1073). Methods commonly employed to determine identity or similarity between two sequences include, but are not limited to, those disclosed in *Guide to HUGO Computers*, Martin J. Bishop, ed., Academic Press, San Diego, 1994, and Carillo, H., and Lipson, D., *SIAM J Applied Math* (1988) 48:1073. Methods to determine identity and similarity are codified in computer programs. Preferred computer programs methods to determine identity and similarity between two sequences include, but are not limited to, GCG program package (Devereux, J., et al., *Nucleic Acids Research* (1984) 12(1):387), BLASTP, BLASTN, and FASTA (Altschul, S. F., et al., *JMolec Biol* (1990) 215:403).

[0077] By way of example, a polynucleotide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:1, that is be 100% identical, or it may include up to a certain integer number of nucleotide alterations as compared to the reference sequence. Such alterations are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion, and wherein said alterations may occur at the 5' or 3' terminal positions of the reference nucleotide sequence or anywhere between those terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. The number of nucleotide alterations is determined by multiplying the total number of nucleotides in SEQ ID NO:1 by the numerical percent of the respective percent identity (divided by 100) and

subtracting that product from said total number of nucleotides in SEQ ID NO:1, or:

$$n_n \leq X_n - (x_n \cdot y),$$

wherein n_n is the number of nucleotide alterations, x_n is the total number of nucleotides in SEQ ID NO:1, and y is 0.50 for 50%, 0.60 for 60%, 0.70 for 70%, 0.80 for 80%, 0.85 for 85%, 0.90 for 90%, 0.95 for 95%, 0.97 for 97% or 1.00 for 100%, and wherein any non-integer product of x_n and y is rounded down to the nearest integer prior to subtracting it from x_n . Alterations of a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may create nonsense, missense or frameshift mutations in this coding sequence and thereby alter the polypeptide encoded by the polynucleotide following such alterations.

[0078] Similarly, a polypeptide sequence of the present invention may be identical to the reference sequence of SEQ ID NO:2, that is be 100% identical, or it may include up to a certain integer number of amino acid alterations as compared to the reference sequence such that the % identity is less than 100%. Such alterations are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion, and wherein said alterations may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between those terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. The number of amino acid alterations for a given % identity is determined by multiplying the total number of amino acids in SEQ ID NO:2 by the numerical percent of the respective percent identity (divided by 100) and then subtracting that product from said total number of amino acids in SEQ ID NO:2, or:

$$n_a \leq x_a - (x_a \cdot y),$$

wherein n_a is the number of amino acid alterations, x_a is the total number of amino acids in SEQ ID NO:2, and y is, for instance 0.70 for 70%, 0.80 for 80%, 0.85 for 85% etc., and wherein any non-integer product of x_a and y is rounded down to the nearest integer prior to subtracting it from x_a . [0079] "Fusion protein" refers to a protein encoded by two, often unrelated, fused genes or fragments thereof. In one example, EP-A-0 464 discloses fusion proteins comprising various portions of constant region of immunoglobulin molecule together with another human protein or part thereof. In many cases, employing an immunoglobulin Fc region as a part of a fusion protein is advantageous for use in therapy and diagnosis resulting in, for example, improved pharmacokinetic properties [see, e.g., EP-A 0 232 262]. On the other hand, for some uses it would be desirable to be able to delete the Fc part after the fusion protein has been expressed, detected and purified. [0080] All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

[illegible]

SEQ ID NO:1

SEQUENCE INFORMATION

[0081]

SEQUENCE LISTING

(1) GENERAL INFORMATION

(1) APPLICANT: Smt thKline Beecham Corporation

(11) TITLE OF THE INVENTION: Sialoadhesin Family Member-2

(SAF-2)

(111) NUMBER OF SEQUENCES: 4

(1v) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Smt thKline Beecham, Corporate Intellectual Property
(B) STREET: Two New Horizons Court
(C) CITY: Brentford

(D) STATE: Middlesex

(E) COUNTRY: United Kingdom

(F) ZIP: TW9 9EP

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Diskette

(B) COMPUTER I BM Compatible

(C) OPERATING SYSTEM: DOS

(D) SOFTWARE: FastSEQ for Windows Version 2.0

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER TO BE ASSIGNED

(B) FILING DATE: 11-MAR-1998

(C) CLASSIFICATION: UNKNOWN

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: 60/041,886

(B) FILING DATE: 02-APR-1997

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: GIDDINGS, Peter John

(B) REGISTRATION NUMBER: 5630

(C) REFERENCE/DOCKET NUMBER: GH-50018

EP 0 924 297 A1

(1X) TELECOMMUNICATION INFORMATION

(A) TELEPHONE: +44 181 975 6331

(B) TELEFAX: +44 181 975 6294

(C) TELETYPE

(2) INFORMATION FOR SEQ ID NO: 1:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2900 base pairs

(B) TYPE: nucleic acid

(C) ST RANDEDNESS: single

(D) TOPOLOGY: Linear

(11) MOLECULE TYPE: CDNA

(41) SEQUENCE DESCRIPTION SEQ ID NO: 1:

55	1500	CAGGACT CCC	TCAT CTGTCT	GACAGCATGG	CAGT CCGCTAC	CCCCCAATTTT	TCCCCAGGCA
	1440	TCTTTGGT AT	GGCCAAAGT C	ACTT CCGT CCGT	CT CCGT CCGT CC	TT CCCACAGG	CCGGCTT CTA
	1380	TT CT CCACAC	CGAGCAT CCA	GCTGGACAC	CT CTGACAGG	ATTGGCCCCC	AGGAT CCGT C
	1320	ACAGGCATGG	AAAGTGCAAA	GGCAT CAGG	GGCT CGGCT	CT CAGGTGAG	TGATGTGGG
50	1260	TT CAT CAT AG	TAGGT CCGT G	CAGGAAGAAA	T CGGCAAGGC	CAGCAGCGGG	CGTGGGGGAT
	1200	CT GGCAGCAG	T CGGGGGAGC	TGGAGCCACA	GCCCTGGCCT	T CCGT CCGT	CTGCAT CAT C
	1140	CT GAGCCT CT	CCCTGCAGAA	TGAGGGGACA	GGCACCCT CAA	GACCTGT AT C	ACAAGTGACA
	1080	AGGAGTGAAG	GGCAATT CAC	CTGGCCGAGCT	CAGAACCGCT C	AGGGCT CCGA	GCACAATTT CC
45	1020	ACCTGTGCCC	CGT CAGCGT C	CT CAAACTCT	GGGTGCTGG	AGCTGCT CG	AGTGACAGTG
	960	CT GGT CTGT G	CTGT CAACAG	CAT CCCCCCT	GCCAGCTGA	GCTGGACCGG	GGGAGGCTG
	900	GCAT CCAACAG	CCCTGGGAAA	TGGT CAT CT	CTTT CAGT CC	TTGAGGGCCA	GT CT CTGCGC
	840	CT CGATGTGT	CTAACCCT CC	TTGGAACCTG	ACCATGACTG	T CTT CCAAGG	AGATGCCACA
40	780	AGCCT CACCT	GT CAGGTGAC	CTTGCCCTGG	ACAGGTGTGA	CCAAGAACCA	TACCGT CCGC
	720	CCCACTACTG	CCCGT CCGT C	AGTGT CACC	CTTAACCCAA	AGCCCCAGGA	CCAAGGCCAC
	660	TGT AAGCAGG	GGAACACCCC	CATGAT CT CC	TGGATTTGGGG	CGT CCGT GT C	CT CCCCAGG
	600	AT CCAAGGA	CCCT AAGAT C	TGGCCACT CC	AGGAACCTGA	CGTGT CT GT	GCCCTGGCC
35	540	TACAAACT A	AGCAGCTGT C	TGTGTTTGTG	ACAGCCCTGA	CCCATAGGC	TGACAT CCGT C
	480	T CAT ATTT CT	TT CGGCT AGA	GAGAGGAAGC	ATGAATGGA	GTTACAAT C	ACAGTTGAA
	420	GACATTTTGA	GCAACGACTG	CT CCGT GAGC	AT CAGAGACG	CCAGGAAGAG	GGAT AAGGGG
	360	ACAAACAACC	CAGACAGAGA	AGTGACAGCA	GAGACCCAGG	GCCGATT CCA	ACT CCGTTGG
30	300	CCAGTT CAT G	GCTACTGGTT	CCGGGACAGA	GACAGACCAT	ACCAAGACCG	TCCAAGTGGCC
	240	CTGTGTCT CC	AGTGCCCGT G	CT CCGT CT CC	TACCCCCAGG	ATGGCTGGAC	TGACT CTGAC
	180	AGACAT AT G	GGATGGTTA	CTTGCTGCAA	GTGACAGAGC	TGGTGAACGT	GACGAGAGGC
25	120	ATGCTGCTGC	TGCTGCTGCT	GCTGCCCCCTG	CT CTGGGGGA	CAAAAGGGAT	GAGGGGAGAC
	60	CCCAACGGCT C	CGCTGAGGAA	CAGACGTT CC	CTGGCGGGCC	TGGCGCGCT C	AAACCCAGAC

Met Leu Leu Leu Leu Leu Leu Pro Leu Leu Trp Gly Thr Lys Gly
1
5
10
15
Met Gly Gly Asp Arg Gln Tyr Gly Asp Gly Tyr Leu Leu Gln Val Gln
20
25
30
Gln Leu Val Thr Val Gln Gln Gly Leu Cys Val His Val Pro Cys Ser
35
40
45

(1) SEQUENCE CHARACTERISTICS
(A) LENGTH: 431 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(11) MOLECULE TYPE: protein
(11) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

(2) INFORMATION FOR SEQ ID NO: 2:

AAAAAAAAAAAAAAAAAAAA
2900
2880
2860
2840
2820
2800
2780
2760
2740
2720
2700
2680
2660
2640
2620
2600
2580
2560
2540
2520
2500
2480
2460
2440
2420
2400
2380
2360
2340
2320
2300
2280
2260
2240
2220
2200
2180
2160
2140
2120
2100
2080
2060
2040
2020
2000
1980
1960
1940
1920
1900
1880
1860
1840
1820
1800
1780
1760
1740
1720
1700
1680
1660
1640
1620
1600
1580
1560

EP 0 924 297 A1

385 390 395 400

Thr Gly Met Glu Asp Ala Lys Ala Ile Arg Gly Ser Ala Ser Gln Val

405 410 415

Ser Asp Val Gly Phe Ser Thr Pro Ser Ile Gln Pro Gly His Leu

420 425 430

(2) INFORMATION FOR SEQ ID NO:3:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 1597 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(11) MOLECULE TYPE: cDNA

(11) SEQUENCE DESCRIPTION: SEQ ID NO:3:

25 GTATGTTGTC TGAATGTG AGCGATAAC AATTTCACAC AGGAACACG TATGACCATG
 120 ATTACGCCAA GCTCTAATAC GACTCACTAT AGGGAAGCT GTACGCCCTG CAGGTACCGT
 180 TCCGGAATTC CCGGGTCCA CCGACGCGTC CGCTGAGGAA CAGACGTTCC CTGGCGGCC
 240 TGGCGCGCTTC AAACCCAGAC ATGCTGCTGC TGTGCTGCT GCTGCCCTG CTCTGGGGGA
 300 CAAAGGGGAT GGAAGGAGAC AGACAAATATG GGGATGGTTA CTGCTGCATA GTGCAGGAGC
 360 TGGTGACCGT GCAGGAGGCG CTGTGTGCTC ATGTGCGCTG CTCCTTCTCC TACCCCCAGG
 420 ATGGCTGGAC TGACTCTGAC CCAGTTCATG GCTACTGGTT CCGGGCAGGA GACAGACCAT
 480 ACCAAGACGC TCCAGTGGCC ACAAAACAAC CAGACAGAGA AGTGCAGGCA GAGACCCACG
 540 GCCGATCCA ACTCCTTGGG GACATTTGGA GCAACGACTG CTCCTGAGC ATCAGAGACG
 600 CCAGGAAGAG GGATAAAGGG TCATATTTCT TTCGCTAGA GAGAGGAAGC ATGAAATGGA
 660 GTTACAATTC ACAAGTTGAAT TACAATACTA AGCAGCTGTC TGTGTTGTG ACAGCCCTGA
 720 CCGTAAAGCC TGAATCTCTC ATCCTAGGGA CCTTAGACTC TGGCCACTCC AGGAACCTGA
 780 CCTGCTCTGT GCCCTGGGCC TGTAAAGCAGG GGAACACCCCC CATGATCTCC TGGATTGGGG
 840 CCTCCTGTCT CCCCCGGGC CCACTAATG CCGCTCTC AGTGCTCACC CTTACCCCAA
 900 AGCCCCCAGGA CCAAGGCAAC AGCCTCACT GTCAGGTGAC CTGCTCGGG ACAGGTGTGA
 960 CCACGAGCACG TACCGTCCG CTCGATGTGT CTTACCCCTCC TTGGAATCTG ACCATGACTG
 1020 TCTTCCAAGG AGATGCCACA GCATCCAACG CCTGGGAAA TGGCTCATCT CTTTCAGTCC
 1080 TTGAAGGGCCA GTCTCTGCG CTCGTCTGTG CTGTCAACAG CAATCCCCCT GCCAAGCTGA
 1140 GCTGGACCCG GGGAGAGCCTG ACCCTGTGCC CCTCAAGCTC CTCAAAACCT GGGCTGCTGG
 1200 AGCTGCCCTCG AGTGCACTG AGGATGAAAG GCGAATCAC CTGCCGAGCT CAGAAACGCTC
 1260 AGGCTCCCA GCACATTTCC CTGAGCCTCT CCTGCAAGAA TGAAGGCGACA GGCACCTCAA
 1320 GACCTGTATC ACAAGTGACA CTGGCAGCAG TCGGGGAGC TGGAGCCACA GCCCTGGCCT
 1380 TCCTGTCTCT CTGCATCATC TTCAATAG TGAAGTCTCT CAGGAAGAAA TCGGCAAGGC
 1440 CAGCAGCCGG CGTGGGGGAT ACAGGCTATGG AAGATGCAAA GGCCATCAGG GGCTCGGCCCT
 1500 CTCAAGGTGAG TGATGTGGGC TTCTCCACAC CGAGCATCCA GCCTGGACAC CTCTGACAGG

ATGCCCCCA GGATCGCTCT GTTGGATG GCCAAGTCA CTTCCTCGTCT CCTCCTCCT
 1560 TCCAGAGCCCG GCTTCTACAG GACTCCCAT CTTCCTG

1597

(2) INFORMATION FOR SEQ ID NO: 4:

(1) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 431 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(11) MOLECULE TYPE: protein

(x1) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

Met Leu Leu Leu Leu Leu Leu Leu Leu Trp Gly Thr Lys Gly
 1 5 10 15
 Met Glu Gly Asp Arg Gln Tyr Gly Asp Gly Tyr Leu Leu Gln Val Gln
 20 25 30 35
 Glu Leu Val Thr Val Gln Gly Leu Cys Val His Val Pro Cys Ser
 40 45 50 55
 Phe Ser Tyr Pro Gln Asp Gly Trp Thr Asp Ser Asp Pro Val His Gly
 60 65 70 75 80
 Tyr Trp Phe Arg Ala Gly Asp Arg Pro Tyr Gln Asp Ala Pro Val Ala
 85 90 95
 Thr Asn Asn Pro Asp Arg Glu Val Gln Ala Glu Thr Gln Gly Arg Phe
 100 105 110
 Asp Ala Arg Lys Arg Asp Lys Gly Ser Tyr Phe Phe Arg Leu Gln Arg
 115 120 125
 Gly Ser Met Lys Trp Ser Tyr Lys Ser Gln Leu Asn Tyr Lys Thr Lys
 130 135 140
 Gln Leu Ser Val Phe Val Thr Ala Leu Thr His Arg Pro Asp Ile Leu
 145 150 155 160
 Ile Leu Gly Thr Leu Glu Ser Gly His Ser Arg Asn Leu Thr Cys Ser
 165 170 175
 Val Pro Trp Ala Cys Lys Gln Gly Thr Pro Pro Met Ile Ser Trp Ile
 180 185 190
 Gly Ala Ser Val Ser Ser Pro Gly Pro Thr Thr Ala Arg Ser Ser Val
 195 200 205
 Leu Thr Leu Thr Pro Lys Pro Gln Asp His Gly Thr Ser Leu Thr Cys

(i) an isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide that has at least

2. An isolated polynucleotide selected from the group consisting of:

(iii) an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:2 or
(iii) an isolated polypeptide which is the amino acid sequence of SEQ ID NO:2

to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2;

(a) 70% identity;
(b) 80% identity;
(c) 90% identity; or
(d) 95% identity

(i) an isolated polypeptide comprising an amino acid sequence selected from the group having at least:

1. An isolated polypeptide selected from the group consisting of:

Claims

210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430

Gln Val Thr Leu Pro Gly Thr Gly Val Thr Thr Ser Thr Val Arg
Leu Asp Val Ser Tyr Pro Pro Trp Asn Leu Thr Met Thr Val Phe Gln
Gly Asp Ala Thr Ala Ser Thr Ala Leu Gly Asn Gly Ser Ser Leu Ser
Val Leu Gln Gly Gln Ser Leu Arg Leu Val Cys Ala Val Asn Ser Asn
Pro Pro Ala Arg Leu Ser Trp Thr Arg Gly Ser Leu Thr Leu Cys Pro
Ser Arg Ser Ser Asn Pro Gly Leu Leu Gln Leu Pro Arg Val His Val
Arg Asp Gln Gly Gln Phe Thr Cys Arg Ala Gln Asn Ala Gln Gly Ser
Gln His Ile Ser Leu Ser Leu Ser Leu Gln Asn Gln Gly Thr Gly Thr
Ser Arg Pro Val Ser Gln Val Thr Leu Ala Val Gly Gly Ala Gly
Ala Thr Ala Leu Ala Phe Leu Ser Phe Cys Ile Ile Phe Ile Ile Val
Arg Ser Cys Arg Lys Lys Ser Ala Arg Pro Ala Ala Gly Val Gly Asp
Thr Gly Met Gln Asp Ala Lys Ala Ile Arg Gly Ser Ala Ser Gln Val
Ser Asp Val Gly Phe Ser Thr Pro Ser Ile Gln Pro Gly His Leu

- 5
 (a) 70% identity;
 (b) 80% identity;
 (c) 90% identity; or
 (d) 95% identity;
- 10
 (a) 70% identity;
 (b) 80% identity;
 (c) 90% identity; or
 (d) 95% identity;
- 15
 (iii) an isolated polynucleotide comprising a nucleotide sequence which has at least:
 over its entire length to a nucleotide sequence encoding the polypeptide of SEQ ID NO:2;
 (a) 70% identity;
 (b) 80% identity;
 (c) 90% identity; or
 (d) 95% identity;
- 20
 (a) 70% identity;
 (b) 80% identity;
 (c) 90% identity; or
 (d) 95% identity;
- 25
 (iv) an isolated polynucleotide comprising a nucleotide sequence encoding the polypeptide of SEQ ID NO:2;
 (vi) an isolated polynucleotide which is the polynucleotide of SEQ ID NO: 1; or
 (vi) an isolated polynucleotide obtainable by screening an appropriate library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or a fragment thereof;
- 30
 3. An antibody immunospecific for the polypeptide of claim 1.
- 35
 4. A method for the treatment of a subject:
 (i) in need of enhanced activity or expression of the polypeptide of claim 1 comprising:
 (a) administering to the subject a therapeutically effective amount of an agonist to said polypeptide; and/or
 (b) providing to the subject an isolated polynucleotide comprising a nucleotide sequence encoding said polypeptide in a form so as to effect production of said polypeptide activity *in vivo*; or
 (ii) having need to inhibit activity or expression of the polypeptide of claim 1 comprising:
 (a) administering to the subject a therapeutically effective amount of an antagonist to said polypeptide; and/or
 (b) administering to the subject a nucleic acid molecule that inhibits the expression of a nucleotide sequence encoding said polypeptide; and/or
 (c) administering to the subject a therapeutically effective amount of a polypeptide that competes with said polypeptide for its ligand, substrate, or receptor.
- 40
 5. A process for diagnosing a disease or a susceptibility to a disease in a subject related to expression or activity of the polypeptide of claim 1 in a subject comprising:
 (a) determining the presence or absence of a mutation in the nucleotide sequence encoding said polypeptide in the genome of said subject; and/or
 (b) analyzing for the presence or amount of said polypeptide expression in a sample derived from said subject.
- 55
 6. A method for screening to identify compounds which stimulate or which inhibit the function of the polypeptide of claim 1 which comprises a method selected from the group consisting of:

(a) measuring the binding of a candidate compound to the polypeptide (or to the cells or membranes bearing the polypeptide) or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound;

(b) measuring the binding of a candidate compound to the polypeptide (or to the cells or membranes bearing the polypeptide) or a fusion protein thereof in the presence of a labeled competitor;

(c) testing whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells or cell membranes bearing the polypeptide;

(d) mixing a candidate compound with a solution containing a polypeptide of claim 1, to form a mixture, measuring activity of the polypeptide in the mixture, and comparing the activity of the mixture to a standard; or

(e) detecting the effect of a candidate compound on the production of mRNA encoding said polypeptide and said polypeptide in cells, using for instance, an ELISA assay.

7. An agonist or an antagonist of the polypeptide of claim 1.

8. An expression system comprising a polynucleotide capable of producing a polypeptide of claim 1 when said expression system is present in a compatible host cell.

9. A process for producing a recombinant host cell comprising transforming or transfecting a cell with the expression system of claim 8 such that the host cell, under appropriate culture conditions, produces a polypeptide comprising an amino acid sequence having at least 70% identity to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2.

10. A recombinant host cell produced by the process of claim 9.

11. A membrane of a recombinant host cell of claim 10 expressing a polypeptide comprising an amino acid sequence having at least 70% identity to the amino acid sequence of SEQ ID NO:2 over the entire length of SEQ ID NO:2.

12. A process for producing a polypeptide comprising culturing a host cell of claim 10 under conditions sufficient for the production of said polypeptide and recovering the polypeptide from the culture.

13. An isolated polynucleotide selected from the group consisting of:

(a) an isolated polynucleotide comprising a nucleotide sequence which has at least 70%, 80%, 90%, 95%, 97% identity to SEQ ID NO:3 over the entire length of SEQ ID NO:3;

(b) an isolated polynucleotide comprising the polynucleotide of SEQ ID NO:3;

(c) the polynucleotide of SEQ ID NO:3; or

(d) an isolated polynucleotide comprising a nucleotide sequence encoding a polypeptide which has at least 70%, 80%, 90%, 95%, 97-99% identity to the amino acid sequence of SEQ ID NO:4 over the entire length of SEQ ID NO:4.

14. A polypeptide selected from the group consisting of:

(a) a polypeptide which comprises an amino acid sequence which has at least 70%, 80%, 90%, 95%, 97-99% identity to that of SEQ ID NO:4 over the entire length of SEQ ID NO:4;

(b) a polypeptide which has an amino acid sequence which is at least 70%, 80%, 90%, 95%, 97-99% identity to the amino acid sequence of SEQ ID NO:4 over the entire length of SEQ ID NO:4;

(c) a polypeptide which comprises the amino acid of SEQ ID NO:4;

(d) a polypeptide which is the polypeptide of SEQ ID NO:4;

(e) a polypeptide which is encoded by a polynucleotide comprising the sequence contained in SEQ ID NO:3.

European Patent Office
 PARTIAL EUROPEAN SEARCH REPORT
 Application Number
 which under Rule 45 of the European Patent Convention EP 98 30 2537 shall be considered, for the purposes of subsequent proceedings, as the European search report



DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
----------	---	-------------------	--

A	SIMMONS D. & SEED B.: "Isolation of a CDNA encoding CD33, a differentiation antigen of myeloid progenitor cells." J. IMMUNOL. vol. 141, 15 October 1988, pages 2797-2800, XP002072092	I-14	C12N15/12 C12N15/85 C12N5/10 C12Q1/68 C07K14/705 C07K16/28 G01N33/50 A61K48/00
---	---	------	---

A	CRÖCKER P.R. ET AL.: "Sialoadhesin, a macrophage sialic acid binding receptor for haemopoietic cells with 17 immunoglobulin-like domains" EMBO J., vol. 13, no. 19, 1994, pages 4490-4503, XP002072093	I-14	TECHNICAL FIELDS SEARCHED (Int.Cl.6) C07K
---	--	------	--

INCOMPLETE SEARCH

The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPC to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.

Claims searched completely :

Claims searched incompletely :

Claims not searched :

Reason for the limitation of the search:

Although claim 4 is directed to a method of treatment of the human/animal body (Article 52(4) EPC), the search has been carried out and based on the alleged effects of the compound/composition.

Place of search	THE HAGUE	Date of completion of the search	22 July 1998	Examiner	Galii, I
-----------------	-----------	----------------------------------	--------------	----------	----------

CATEGORY OF CITED DOCUMENTS

- 1 : theory or principle underlying the invention
 E : earlier patent document, but published on, or after the filing date
 D : document cited in the application
 L : document cited for other reasons
 A : technological background
 O : non-written disclosure
 P : intermediate document
 X : particularly relevant if taken alone
 Y : particularly relevant if combined with another document of the same category
 Z : document of the same category
 A : technological background
 O : non-written disclosure
 P : intermediate document

EPC FORM 1503 03 92 (P4/C07)

DOCUMENTS CONSIDERED TO BE RELEVANT		CLASSIFICATION OF THE APPLICATION (INCL.6)	
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
A	KELM S ET AL: "THE SIALOADHESINS - A FAMILY OF SIALIC ACID-DEPENDENT CELLULAR RECOGNITION MOLECULES WITHIN THE IMMUNOGLOBULIN SUPERFAMILY" GLYCOCOMJUGATE JOURNAL, vol. 13, no. 6, 1996, pages 913-926, XP002066896 * the whole document *	1-14	
A	ZANNETTINO A C W ET AL: "A POWERFUL NEW TECHNIQUE FOR ISOLATING GENES ENCODING CELL SURFACE ANTIGENS USING RETROVIRAL EXPRESSION CLONING" JOURNAL OF IMMUNOLOGY, vol. 156, no. 2, 15 January 1996, pages 611-620, XP002066893 * abstract *	1-14	TECHNICAL FIELDS SEARCHED (INCL.6)
A	ADAMS M D ET AL: "COMPLEMENTARY DNA SEQUENCING: EXPRESSED SEQUENCE TAGS AND HUMAN GENOME PROJECT" SCIENCE, vol. 252, no. 5013, 21 June 1991, pages 1651-1656, XP000645049 * abstract *	1-14	
P, X	DATABASE GENBANK Accession No. AA683022, 10 December 1997 HILLIER L ET AL: "EST; Homo sapiens clone 970578, contains OFR.b1 repetitive element." XP002072094 Compare nt 2859-2519 of Seq. ID 3 with AA683022	1,2,13, 14	

EPO FORM 1501 03.82 (PACT/0)



